

MYOCARDIAL ISCHEMIA AND INFARCTION

THE PROGNOSTIC VALUE OF THE THROMBOLYSIS IN MYOCARDIAL INFARCTION (TIMI) RISK INDEX AFTER PRIMARY PERCUTANEOUS CORONARY INTERVENTION FOR ST-ELEVATION MYOCARDIAL INFARCTION AT LONG-TERM FOLLOW-UP

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Monday, April 04, 2011, 9:30 a.m.-10:45 a.m.

Session Title: Management of ACS

Abstract Category: 4. Unstable Ischemic Syndrome/Long-Term Outcome

Session-Poster Board Number: 1073-371

Authors: Pier Woudstra, Peter Damman, Karel T. Koch, José P.S. Henriques, Jan Baan, Marije M. Vis, Jan J. Piek, Jan G.P. Tijssen, Robbert J. de Winter, Department of Cardiology, Academic Medical Centre, University of Amsterdam, Amsterdam, The Netherlands

Background: In order to optimize the use of novel therapies and improve outcomes in high-risk ST-elevation myocardial infarction (STEMI) patients, risk stratification is of importance. The Thrombolysis In Myocardial Infarction Risk Index (TIMI-RI) for STEMI was developed to get insight in short-term outcomes. In this study we assessed discriminative value of the TIMI-RI in predicting short and long-term mortality in STEMI patients treated with primary percutaneous coronary intervention (PPCI).

Methods: We used registry data from STEMI patients undergoing PPCI in a high-volume PCI centre in The Netherlands. Data on all-cause mortality was collected up to 5 years of follow-up. Patients were divided into tertiles based on TIMI-RI (<17.4, >17.4 and <27.9, >27.9). Cumulative event rates were estimated with the Kaplan-Meier method and compared with a log-rank test. A landmark was set at 30 days. Discrimination of the TIMI-RI was assessed with C-statistics.

Results: We included 1806 patients with a mean follow-up of 1046 days. The cumulative mortality is shown in figure 1. Mortality was significantly higher in higher TIMI-RI categories before and after the 30-day landmark ($p < 0.001$). Overall the discriminative value of the TIMI-RI for mortality estimated by the C statistic was 0.791. The C statistic of the TIMI-RI was respectively 0.851 and 0.720 for short and long-term prediction of mortality.

Conclusion: TIMI-RI predicts not only short but also long-term mortality in STEMI patients treated with PPCI.

Figure 1. Landmark cumulative mortality curves according to TIMI risk index

